



Expert opinion on evidence after the 2020 Korean Cardiopulmonary Resuscitation Guidelines: a secondary publication

Sung Phil Chung¹, Youdong Sohn², Jisook Lee³, Youngsuk Cho²,
Kyoung-Chul Cha⁴, Ju Sun Heo^{5,6}, Ai-Rhan Ellen Kim⁷, Jae Guk Kim⁸,
Han-Suk Kim⁹, Hyungoo Shin¹⁰, Chiwon Ahn¹¹, Ho Geol Woo¹²,
Byung Kook Lee¹³, Yong Soo Jang⁸, Yu Hyeon Choi¹⁴, Sung Oh Hwang⁴;
on behalf of the Guideline Committee of the Korean Association of
Cardiopulmonary Resuscitation (KACPR)

- ¹Department of Emergency Medicine, Yonsei University College of Medicine, Seoul, Korea
²Department of Emergency Medicine, Hallym University Kangdong Sacred Heart Hospital, Seoul, Korea
³Department of Emergency Medicine, Ajou University School of Medicine, Suwon, Korea
⁴Department of Emergency Medicine, Yonsei University Wonju College of Medicine, Wonju, Korea
⁵Department of Pediatrics, Korea University College of Medicine, Seoul, Korea
⁶Institute of Nano, Regeneration, Reconstruction, Korea University, Seoul, Korea
⁷Department of Pediatrics, CHA University Ilsan Medical Center, Goyang, Korea
⁸Department of Emergency Medicine, Hallym University Kangnam Sacred Heart Hospital, Seoul, Korea
⁹Department of Pediatrics, Seoul National University College of Medicine, Seoul, Korea
¹⁰Department of Emergency Medicine, Hanyang University College of Medicine, Seoul, Korea
¹¹Department of Emergency Medicine, Chung-Ang University College of Medicine, Seoul, Korea
¹²Department of Neurology, Kyung Hee University College of Medicine, Seoul, Korea
¹³Department of Emergency Medicine, Chonnam National University Medical School, Gwangju, Korea
¹⁴Department of Pediatrics, Seoul Medical Center, Seoul, Korea

Considerable evidence has been published since the 2020 Korean Cardiopulmonary Resuscitation Guidelines were reported. The International Liaison Committee on Resuscitation (ILCOR) also publishes the Consensus on CPR and Emergency Cardiovascular Care Science with Treatment Recommendations (CoSTR) summary annually. This review provides expert opinions by reviewing the recent evidence on CPR and ILCOR treatment recommendations. The authors reviewed the CoSTR summary published by ILCOR in 2021 and 2022. PICO (patient, intervention, comparison, outcome) questions for each topic were reviewed using a systemic or scoping review methodology. Two experts were appointed for each question and reviewed the topic independently. Topics suggested by the reviewers for revision or additional description of the guidelines were discussed at a consensus conference. Forty-three questions were reviewed, including 15 on basic life support, seven on advanced life support, two on pediatric life support, 11 on neonatal life support, six on education and teams, one on first aid, and one related to COVID-19. Finally, the current Korean CPR Guideline was maintained for 28 questions, and expert opinions were suggested for 15 questions.

Keywords Heart arrest; Cardiopulmonary resuscitation; Expert opinion

Received: 2 August 2023
Revised: 18 August 2023
Accepted: 20 August 2023

Correspondence to: Sung Oh Hwang
Department of Emergency Medicine,
Yonsei University Wonju College of
Medicine, 20 Ilsan-ro, Wonju 26426,
Korea
Email: shwang@yonsei.ac.kr

This article is a translated secondary publication of "Expert opinion on evidence after 2020 Korean Cardiopulmonary Resuscitation Guidelines," published in the *Journal of the Korean Society of Emergency Medicine* 2023;34(4):287-96 (in Korean).



How to cite this article:

Chung SP, Sohn Y, Lee J, Cho Y, Cha KC, Heo JS, Kim ARE, Kim JG, Kim HS, Shin H, Ahn C, Woo HG, Lee BK, Jang YS, Choi YH, Hwang SO; on behalf of the Guideline Committee of the Korean Association of Cardiopulmonary Resuscitation (KACPR). Expert opinion on evidence after the 2020 Korean Cardiopulmonary Resuscitation Guidelines: a secondary publication. *Clin Exp Emerg Med* 2023;10(4):382-392. <https://doi.org/10.15441/ceem.23.102>

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0/>).

Capsule Summary

What is already known

Cardiopulmonary resuscitation (CPR) guidelines are revised every 5 years. Since the 2020 guidelines were published, the International Liaison Committee on Resuscitation (ILCOR) has published evidence summaries every year. It is necessary to update the recommendations continuously, reflecting the new evidence.

What is new in the current study

This review summarizes the expert opinions on the new evidence since the 2020 Korean Cardiopulmonary Resuscitation Guidelines. Forty-three PICO (patient, intervention, comparison, outcome) questions were reviewed, and consensus opinions were suggested for 15 questions.

INTRODUCTION

The Korean cardiopulmonary resuscitation (CPR) guidelines have been updated periodically since they were first developed in 2006, with the fourth revised version published in 2020 [1]. The International Liaison Committee on Resuscitation (ILCOR) reviews the latest evidence and publishes the Consensus on CPR and Emergency Cardiovascular Care Science with Treatment Recommendation (CoSTR) every 5 years starting in 2005, which is summarized as a practice guideline. The Korean CPR guidelines can be developed based on the CoSTR because the Korean Association of Cardiopulmonary Resuscitation (KACPR) participated in the evidence review process as a member of the Resuscitation Council of Asia, a member organization of the ILCOR [2].

As research and publications in the field of resuscitation have increased, the need to shorten the update cycle of CPR guidelines has increased. Accordingly, the ILCOR has been conducting evidence reviews and publishing CoSTR summaries annually since 2017 [3]. It also releases an annual update of CoSTR summaries since the introduction of the 2020 guidelines [4,5], which can be found on its website (costr.ilcor.org). The Guideline Committee of KACPR concluded that it would be better to review recent evidence and to update the necessary contents before the next revision of the Korean CPR guidelines, which is scheduled for 2025. This review includes expert consensus opinions regarding the evidence on CPR published after the 2020 Korean CPR Guidelines.

EVIDENCE REVIEW METHODOLOGY

The evidence review focused on the topics of the CoSTR summaries published by the ILCOR in 2021 [4] and 2022 [5]. The members of the CPR Guidelines Committee and the evidence reviewers included experts recommended by eight professional organiza-

tions related to the CPR guidelines. The committee members and evidence reviewers were experts with experience using the methodology for revising the guidelines, including literature search, systemic review and meta-analysis, and the GRADE (Grading of Recommendations Assessment, Development, and Evaluation) method. Members of the CPR Guidelines Committee selected items that required evidence review among the topics of CoSTR summaries. For review of the topics, the PICO (patient, intervention, comparison, outcome) format was used. For the evidence review, domestic papers as well as papers published in international journals were reviewed. PubMed, Embase, Cochrane Central Register of Controlled Trials (CENTRAL), Education Resources Information Center (ERIC), and KoreaMed were utilized for literature search. Since the 2020 Korean CPR Guidelines included papers published up to September 2020, papers published from October 2020 to May 2023 were included in this review. For topics not covered in the 2020 Korean CPR Guidelines, papers prior to 2020 were also reviewed. The review process used one of three methodologies: systematic review, scoping review, or evidence update. Among the PICO questions on first aid, those unrelated to CPR were excluded. A total of 43 PICO questions was selected for the review, including 15 on basic life support, seven on advanced life support, two on pediatric life support, 11 on neonatal life support, six on education implementation team, one on first aid, and one on COVID-19.

Two experts independently reviewed each PICO question. If one reviewer suggested the need for revision or addition of the 2020 Korean CPR Guidelines, the PICO question was sent to the consensus workshop for discussion and decision (Fig. 1). As a result, new expert consensus opinions were presented for 15 PICO questions (Table 1).




Fig. 1. The process of guideline update. ILCOR, International Liaison Committee on Resuscitation; CoSTR, Consensus on CPR and Emergency Cardiovascular Care Science with Treatment Recommendations.

BASIC LIFE SUPPORT

Video-based dispatch system

Several studies have compared CPR quality to evaluate the effectiveness of video-based emergency dispatch systems. However, only two retrospective observational studies reported in Korea [6,7] compared patient survival as an outcome. The implementation of video-based emergency dispatch systems compared to that of the conventional audio-based systems improved the survival-to-discharge rate (22.3% vs. 10.7%; odds ratio [OR], 2.33; $P < 0.001$) and the rate of good neurological outcomes at discharge (16.0% vs. 6.3%; OR, 2.77; $P < 0.001$) [8]. The 2020 Korean CPR Guidelines recommend dispatcher-assisted CPR; however, only the use of speaker phones or hands-free functions is recommended during dispatcher-assisted CPR. We recommend implementing a video-based dispatch system for improving the survival rate and neurological outcomes of patients with cardiac arrest.

CPR during transport

In an observational study reported from North America [9], the intra-arrest transport group compared with the on-site CPR group had a lower survival-to-discharge rate (4.0% vs. 8.5%; risk difference, 4.6%; range, 4.0%–5.1%) and poor neurological outcomes (2.9% vs. 7.1%; risk difference, 4.2%; range, 3.5%–4.9%). However, it is difficult to generalize this result because the prehospital emergency medical services systems in North America differ from those in Korea in terms of the composition of emergency medical personnel, level of prehospital emergency care, and related legal

regulations. The ILCOR suggested that CPR should be performed on-site, except in cases where the need for transport is clear, such as in cases of extracorporeal membrane oxygenation (ECMO) candidates.

The 2020 Korean CPR Guidelines recommend that transfer to the hospital be considered if spontaneous circulation is not restored after 6 minutes of basic life support or 10 minutes of advanced life support at the scene. For CPR during transport, the Guideline Committee decided to maintain current recommendations. During transport to a hospital, high-quality CPR should be maintained.

In-water resuscitation for individuals who experienced drowning

In a retrospective observational study [10] comparing a group that received rescue ventilation for 1 minute in water when drowning and another that did not, initial survival (94.7% vs. 37.0%, $P < 0.001$), survival-to-discharge (87.5% vs. 25%, $P = 0.005$), and good neurological outcome (52.6% vs. 7.4%, $P = 0.001$) rates were higher in the group that received rescue ventilation. In-water resuscitation can be considered in situations where adequately trained personnel can use appropriate equipment.

ECMO for individuals who experienced drowning

In the 2020 Korean CPR Guidelines, there was no recommendation for the use of ECMO in individuals who experienced drowning. The ILCOR analyzed two retrospective observational studies [11,12] and 11 case reports and stated that ECMO can be consid-

Table 1. Topics reviewed by evidence reviewers

Topic	Recommendation summary
Basic life support	
Video-based dispatch system	We recommend implementing a video-based dispatch system.
CPR during transport	We recommend that transfer to the hospital be considered if spontaneous circulation is not restored after 6 min of basic life support or 10 min of advanced life support at the scene.
In-water resuscitation in drowning	In-water resuscitation can be considered in situations where adequately trained personnel can use appropriate equipment.
ECMO in drowning	ECMO may be beneficial in patients with cardiac arrest due to drowning or in patients with severe respiratory failure due to drowning.
C-A-B or A-B-C in drowning	The C-A-B sequence, starting with chest compressions, is recommended for the sequence of CPR even for a drowning victim.
Advanced life support	
Targeted temperature management after cardiac arrest	We recommend that the target temperature should be maintained at 32–36 °C for at least 24 hr, as recommended by the 2020 Korean CPR Guidelines.
Vasopressin and corticosteroids for cardiac arrest	We suggest considering using vasopressin and steroid combination therapy during CPR in an IHCA setting.
Consciousness during CPR	Sedatives, analgesics, or both may be used in very small doses, if possible, to reduce pain and suffering in conscious patients during CPR. Neuromuscular blocking drugs should not be administered alone to conscious patients.
CPR and defibrillation in the prone position	CPR can be started in patients in the prone position and a secure airway if the change to the supine position is not possible.
Pediatric life support	
PEWS with or without rapid response teams	A scoring system that can detect early warning signs of pediatric cardiac arrest considering the available resources of each hospital is recommended. No recommendation on the pediatric rapid response team.
Neonatal life support	
Cord management at birth for preterm infants	We suggest delayed cord clamping of 30 sec or longer in preterm infants less than 34 wk of gestation who do not require resuscitation, and intact cord milking as a reasonable alternative to delayed cord clamping in preterm infants between 28 and 33 wk of gestation.
Education	
Pre-arrest prediction of survival following IHCA	The use of these predictive scales in patients with IHCA is not recommended.
Basic life support training for potential rescuers for the population at a high risk of cardiac arrest	We recommend that basic life support should be taught to potential rescuers of the population at high risk of cardiac arrest.
Blended learning for life support education	We recommend developing and implementing a blended form of CPR education in addition to the conventional training methods.
Faculty development approaches	We recommend introducing the instructor training program.

CPR, cardiopulmonary resuscitation; ECMO, extracorporeal membrane oxygenation; C-A-B, circulation-airway-breathing; A-B-C, airway-breathing-circulation; IHCA, in-hospital cardiac arrest; PEWS, Pediatric Early Warning Score.

ered in some patients with cardiac arrest who do not respond to conventional CPR. Therefore, ECMO may be beneficial in patients with cardiac arrest or severe respiratory failure due to drowning.

The C-A-B or A-B-C approach in cases of drowning

Nine studies were reviewed; however, none compared the order of CPR in cases of drowning [10,13,14]. The ILCOR suggested, with expert consensus, starting CPR with chest compressions (the C [circulation]-A [airway]-B [breathing] approach) for the layperson, whereas healthcare providers could consider starting rescue ventilation before chest compressions (the A-B-C approach). The 2020 Korean Guidelines do not include recommendations on whether the CPR sequence should be changed depending on circumstances, including drowning. In addition to the suggestion of IL-

COR, considering the simplicity and practicality of CPR training, the C-A-B sequence, starting with chest compressions, is recommended for the sequence of CPR even for a drowning victim.

ADVANCED LIFE SUPPORT

TTM after cardiac arrest

In a targeted temperature management (TTM) trial [15] comparing target temperatures of 33 °C versus 36 °C and the TTM2 trial [16] comparing 33 °C versus normothermia below 37.5 °C (fever prevention), no difference was observed in survival to discharge, 180-day survival, or neurological outcomes at 180 days. According to a network meta-analysis [17] comparing the effects of target temperatures in 10 randomized controlled studies, compared

to normothermia (37–37.8 °C), body temperatures of 31–32, 33–34, and 35–36 °C did not improve survival or neurological outcomes. The ILCOR recommended maintaining the body temperature below 37.5 °C because it can reduce the burden on medical personnel and side effects and the use of fever prevention instead of maintaining normothermia [18]. In the TTM2 study, more than 90% of the cardiac arrests were witnessed, and 62% to 63% of initial rhythms were ventricular fibrillation. The characteristics of out-of-hospital cardiac arrest (OHCA) patients in Korea are different from those of patients included in the TTM and TTM2 studies. Compared to patients enrolled in the TTM and TTM2 studies, Korean OHCA patients had a much lower incidence of a shockable rhythm and a longer time from collapse to recovery of spontaneous circulation, which might be associated with a higher chance of a severe post-cardiac arrest syndrome. Therefore, it is necessary to consider differences in the characteristics of cardiac arrest patients in Korea. In the fever prevention group of the TTM2 study [16], acetaminophen and a method of removing clothes and lowering the room temperature were used; temperature control devices were used to control body temperature in 46% of cases. Retrospective studies have suggested that hypothermia can improve neurological outcomes in patients with severe cerebral ischemic injury [19–21]. We recommend that target temperature should be maintained at 32–36 °C for at least 24 hours, as recommended by the 2020 Korean CPR Guidelines.

Vasopressin and corticosteroids for cardiac arrest

Three randomized controlled trials [22–24] including patients with in-hospital cardiac arrest showed that the use of vasopressin and steroids did not improve the survival to discharge rate (OR, 1.39; 95% confidence interval [CI], 0.90–2.14) or rate of good neurological outcomes (OR, 1.64; 95% CI, 0.99–2.72). However, it increased the rate of return of spontaneous circulation (ROSC; OR, 2.09; 95% CI, 1.54–2.84). The ILCOR recommended not using the vasopressin and steroid combination therapy during CPR because it has not been associated with any significant difference in survival, and no study including patients with OHCA has been conducted. The 2020 Korean Guidelines did not provide any recommendations on the combined use of vasopressin and steroids during CPR. Considering that the combination of these drugs increases the ROSC rate and that these drugs are commonly used in hospitals, we suggest vasopressin and steroid combination therapy during CPR in an in-hospital cardiac arrest setting.

Consciousness during CPR

There have been three observational studies; one cross-sectional study; and several case reports on pain, anxiety, agitation, and

posttraumatic stress disorder in conscious patients during CPR. The studies reported that they verbally reassured the patients or administered sedatives or neuromuscular blockers [25–49]. Following the results of the scoping review conducted by the ILCOR, we recommend the following: (1) sedatives, analgesics, or both may be used in very small doses, if possible, to reduce pain and suffering in conscious patients during CPR; (2) neuromuscular blocking drugs should not be administered alone to conscious patients; (3) the optimal drug regimen for sedation and analgesia during CPR is unclear, and a regimen commonly used in critically ill patients can be used.

CPR and defibrillation in the prone position

Twenty adult and 12 pediatric cases of cardiac arrest in the prone position have been reported. Most of these cases were observed in the operating room and one case in the intensive care unit [50–68]. There was no significant difference in the rate of ROSC or survival discharge among patients for whom chest compressions were started immediately in the prone position compared with patients for which chest compressions were started after changing the position from prone to supine. Arterial pressure during CPR was higher in the prone position group [69,70]. The end-tidal carbon dioxide partial pressure in five adults and two children was 10 mmHg or higher, and the time interval from collapse to defibrillation was shortened when defibrillation was performed in the prone position [54–56,61,65,71–73]. As recommended by ILCOR, we recommend the following for adults with cardiac arrest in the prone position: (1) if cardiac arrest occurs in the prone position with a secured airway, CPR can be started if change to the supine position is not possible or poses a significant risk to the patient; (2) if cardiac arrest occurs in the prone position without a secure airway, the patient position should be changed to supine and CPR should be initiated as soon as possible; (3) if a patient with cardiac arrest is in the prone position and cannot immediately be placed in the supine position, defibrillation can be attempted in the prone position.

PEDIATRIC LIFE SUPPORT

Pediatric early warning scores and pediatric rapid response teams

One randomized controlled trial [74] and 11 cohort studies [75–85] of pediatric early warning scoring systems or implementation of a pediatric rapid response system were reviewed. The use of the pediatric early warning score tended to reduce the incidence of in-hospital cardiac arrest, mortality, and unexpected clinical deterioration. However, since research on the elements to be in-

cluded in the pediatric early warning scoring system is lacking, a scoring system that can detect early warning signs of pediatric cardiac arrest, considering the available resources of each hospital, is recommended. Operation of a pediatric rapid response team is associated with a considerable decline in the preintervention trajectory of critical deterioration and a decreased likelihood of respiratory and cardiopulmonary arrest outside of the critical care unit. However, considering that the medical resources and hospital environment of the hospital where the study was conducted were different from those in Korea, the experts decided to make recommendations on the pediatric rapid response team after additional research results in Korea are released.

NEONATAL LIFE SUPPORT

Cord management at birth for preterm infants

According to a systematic review [86] comparing delayed cord clamping and early cord clamping in preterm infants with a gestational age less than 34 weeks, delayed cord clamping resulted in significantly higher hemoglobin and hematocrit within 24 hours after birth and hematocrit at 7 days after birth and lowest mean arterial pressure within 12 hours after birth. Furthermore, the risk of using inotropics due to hypotension and blood transfusions within 24 hours after birth was significantly lower in these infants. When intact cord milking and early cord clamping were compared, hemoglobin and hematocrit levels within 24 hours after birth were significantly higher and the risk of using inotropics due to hypotension and blood transfusions within 24 hours after birth was significantly lower in infants with intact cord milking. On the other hand, comparison of delayed cord clamping and intact cord milking revealed no significant intergroup differences in these parameters. Based on these findings, the 2021 ILCOR CoSTR suggests delayed cord clamping for more than 30 seconds in preterm infants less than 34 weeks of gestational age who do not require immediate resuscitation and intact cord milking as a reasonable alternative to delayed cord clamping in preterm infants between 28 and 33 weeks of gestation. The 2020 Korean CPR Guidelines recommend that umbilical cord milking not be performed in infants less than 28 weeks of gestation due to the increased risk of intraventricular hemorrhage.

We suggest delayed cord clamping of 30 seconds or longer in preterm infants less than 34 weeks of gestation who do not require resuscitation and intact cord milking as a reasonable alternative to delayed cord clamping in preterm infants between 28 and 33 weeks of gestation.

EDUCATION, IMPLEMENTATION, AND TEAM

Pre-arrest prediction of survival following in-hospital cardiac arrest

Pre-arrest clinical prediction rules, such as pre-arrest morbidity, prognosis after resuscitation, and good outcome following attempted resuscitation scores, have been studied for predicting the prognosis of patients with in-hospital cardiac arrest; however, reliable results have not been achieved [87–93]. Therefore, the use of these predictive scales in patients with in-hospital cardiac arrest is not recommended. In addition, since there have been no related studies in pediatric patients, no recommendations can be made regarding the use of prognostic predictive scales for children with in-hospital cardiac arrest.

Basic life support training for potential rescuers of populations at high risk of cardiac arrest

Potential rescuers, such as family members of high-risk patients, are less likely to voluntarily participate in CPR training but are willing to receive training [94,95]. Several studies [94–99] have recommended that basic life support be taught to potential rescuers of high-risk patients for OHCA, and that emergency staff should encourage potential rescuers to participate in basic life support. We recommend that basic life support be taught to potential rescuers of the population at high risk of cardiac arrest.

Blended learning for life support education

Blended learning is an educational method that combines face-to-face and non-face-to-face forms and was introduced in the 2020 Korean CPR Guidelines. Following the COVID-19 pandemic and the development of information technology, the use of various non-face-to-face methods has become more common in medical education [100–102]. Therefore, if resources and conditions allow, it is recommended to develop and implement a blended form of CPR education in addition to the conventional training methods.

Faculty development approach

Continuing CPR education for the public and emergency medical providers is important to increase survival rates after cardiac arrest. To provide continuous CPR training, an instructor training curriculum to teach trainees is important. Although many studies have explained the necessity of CPR instructor training programs, no study has reported that patient outcomes improved with the introduction of the instructor training program [103,104]. Nevertheless, because the instructor training program is an important factor in the teaching method and performance of trainees, it should be introduced.

CONCLUSION

The Korean CPR guidelines are revised every 5 years. Considering the situation in which new evidence continues to be published, it is necessary to update CPR guidelines to reflect the latest evidence between revision cycles. This review summarizes expert opinions based on the CoSTR summary published by the ILCOR since the publication of the 2020 Korean CPR Guidelines. We hope that this review will contribute to improving the survival of patients with cardiac arrest.

ETHICS STATEMENT

Not applicable.

CONFLICT OF INTEREST

No potential conflict of interest relevant to this article was reported.

FUNDING

None.

ACKNOWLEDGMENTS

The authors thank the following members for their role in the Guideline Committee of Korean Association of Cardiopulmonary Resuscitation (KACPR): Do Kyun Kim (Seoul National University Hospital, Seoul, Korea), Jin Tae Kim (Seoul National University Hospital), Mi Jin Lee (Kyungpook National University, Daegu, Korea), Joo Young Lee (Seoul St. Mary's Hospital, College of Medicine, The Catholic University of Korea, Seoul, Korea), Myung Ja Cho (Republic of Korea National Red Cross, Wonju, Korea), Eun Sun Jin (Kyunghee University Hospital at Gangdong, Seoul, Korea), and Seung Tae Han (Republic of Korea Special Warfare School, Seoul, Korea).

AUTHOR CONTRIBUTIONS

Conceptualization: SOH, SPC; Methodology: SOH, SPC; Project administration: SPC; Supervision: SPC; Writing-original draft: YS, JL, YC, KCC, JSH, AREK, JGK, HSK, HS, CA, HGW, BKL, YSJ, YHC; Writing-review & editing: SOH. All authors read and approved the final manuscript.

ORCID

Sung Phil Chung	https://orcid.org/0000-0002-3074-011X
Youdong Sohn	https://orcid.org/0000-0001-8789-0090
Jisook Lee	https://orcid.org/0000-0002-0522-1350
Youngsuk Cho	https://orcid.org/0000-0002-7648-924X
Kyoung-Chul Cha	https://orcid.org/0000-0003-1818-2466
Ju Sun Heo	https://orcid.org/0000-0001-8928-289X
Ai-Rhan Ellen Kim	https://orcid.org/0000-0002-9859-3021
Jae Guk Kim	https://orcid.org/0000-0002-2446-4497
Han-Suk Kim	https://orcid.org/0000-0002-9777-3231
Hyungoo Shin	https://orcid.org/0000-0003-3203-1672
Chiwon Ahn	https://orcid.org/0000-0002-1813-1098
Ho Geol Woo	https://orcid.org/0000-0001-6489-0100
Byung Kook Lee	https://orcid.org/0000-0003-3571-9448
Yong Soo Jang	https://orcid.org/0000-0001-5964-1580
Yu Hyeon Choi	https://orcid.org/0000-0002-3057-0886
Sung Oh Hwang	https://orcid.org/0000-0003-4585-3181

REFERENCES

1. Hwang SO, Cha KC, Jung WJ, et al. 2020 Korean guidelines for cardiopulmonary resuscitation. Part 1. Update process and highlights. *Clin Exp Emerg Med* 2021;8(S):S1-7.
2. Perkins GD, Neumar R, Monsieurs KG, et al. The International Liaison Committee on Resuscitation: review of the last 25 years and vision for the future. *Resuscitation* 2017;121:104-16.
3. Morley PT. Towards a more continuous evidence evaluation: a collaborative approach to review the resuscitation science. *Resuscitation* 2017;118:A1-2.
4. Wyckoff MH, Singletary EM, Soar J, et al. 2021 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science with Treatment Recommendations: summary from the basic life support; advanced life support; neonatal life support; education, implementation, and teams; first aid task forces; and the COVID-19 Working Group. *Resuscitation* 2021;169:229-311.
5. Wyckoff MH, Greif R, Morley PT, et al. 2022 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science with Treatment Recommendations: summary from the basic life support; advanced life support; pediatric life support; neonatal life support; education, implementation, and teams; and first aid task forces. *Circulation* 2022;146:e483-557.
6. Lee HS, You K, Jeon JP, Kim C, Kim S. The effect of video-instructed versus audio-instructed dispatcher-assisted cardio-

- pulmonary resuscitation on patient outcomes following out of hospital cardiac arrest in Seoul. *Sci Rep* 2021;11:15555.
7. Lee SY, Song KJ, Shin SD, Hong KJ, Kim TH. Comparison of the effects of audio-instructed and video-instructed dispatcher-assisted cardiopulmonary resuscitation on resuscitation outcomes after out-of-hospital cardiac arrest. *Resuscitation* 2020;147:12–20.
 8. Bielski K, Bottiger BW, Pruc M, et al. Outcomes of audio-instructed and video-instructed dispatcher-assisted cardiopulmonary resuscitation: a systematic review and meta-analysis. *Ann Med* 2022;54:464–71.
 9. Grunau B, Kime N, Leroux B, et al. Association of intra-arrest transport vs continued on-scene resuscitation with survival to hospital discharge among patients with out-of-hospital cardiac arrest. *JAMA* 2020;324:1058–67.
 10. Szpilman D, Soares M. In-water resuscitation: is it worthwhile? *Resuscitation* 2004;63:25–31.
 11. Burke CR, Chan T, Brogan TV, et al. Extracorporeal life support for victims of drowning. *Resuscitation* 2016;104:19–23.
 12. Champigneulle B, Bellenfant-Zegdi F, Follin A, et al. Extracorporeal life support (ECLS) for refractory cardiac arrest after drowning: an 11-year experience. *Resuscitation* 2015;88:126–31.
 13. Bierens J, Abelairas-Gomez C, Barcala Furelos R, et al. Resuscitation and emergency care in drowning: a scoping review. *Resuscitation* 2021;162:205–17.
 14. Mtaweh H, Kochanek PM, Carcillo JA, Bell MJ, Fink EL. Patterns of multiorgan dysfunction after pediatric drowning. *Resuscitation* 2015;90:91–6.
 15. Nielsen N, Wetterslev J, Cronberg T, et al. Targeted temperature management at 33°C versus 36°C after cardiac arrest. *N Engl J Med* 2013;369:2197–206.
 16. Dankiewicz J, Cronberg T, Lilja G, et al. Hypothermia versus normothermia after out-of-hospital cardiac arrest. *N Engl J Med* 2021;384:2283–94.
 17. Fernando SM, Di Santo P, Sadeghirad B, et al. Targeted temperature management following out-of-hospital cardiac arrest: a systematic review and network meta-analysis of temperature targets. *Intensive Care Med* 2021;47:1078–88.
 18. Sandroni C, Natalini D, Nolan JP. Temperature control after cardiac arrest. *Crit Care* 2022;26:361.
 19. Callaway CW, Coppler PJ, Faro J, et al. Association of initial illness severity and outcomes after cardiac arrest with targeted temperature management at 36 °C or 33 °C. *JAMA Netw Open* 2020;3:e208215.
 20. Nishikimi M, Ogura T, Nishida K, et al. Outcome related to level of targeted temperature management in postcardiac arrest syndrome of low, moderate, and high severities: a Nationwide Multicenter Prospective Registry. *Crit Care Med* 2021;49:e741–50.
 21. Nutma S, Tjepkema-Cloostermans MC, Ruijter BJ, et al. Effects of targeted temperature management at 33 °C vs. 36 °C on comatose patients after cardiac arrest stratified by the severity of encephalopathy. *Resuscitation* 2022;173:147–53.
 22. Andersen LW, Isbye D, Kjærgaard J, et al. Effect of vasopressin and methylprednisolone vs placebo on return of spontaneous circulation in patients with in-hospital cardiac arrest: a randomized clinical trial. *JAMA* 2021;326:1586–94.
 23. Mentzelopoulos SD, Malachias S, Chamos C, et al. Vasopressin, steroids, and epinephrine and neurologically favorable survival after in-hospital cardiac arrest: a randomized clinical trial. *JAMA* 2013;310:270–9.
 24. Mentzelopoulos SD, Zakynthinos SG, Tzoufi M, et al. Vasopressin, epinephrine, and corticosteroids for in-hospital cardiac arrest. *Arch Intern Med* 2009;169:15–24.
 25. Gamper G, Willeit M, Sterz F, et al. Life after death: posttraumatic stress disorder in survivors of cardiac arrest: prevalence, associated factors, and the influence of sedation and analgesia. *Crit Care Med* 2004;32:378–83.
 26. Parnia S, Spearpoint K, de Vos G, et al. AWARE-Awareness during Resuscitation: a prospective study. *Resuscitation* 2014;85:1799–805.
 27. Olaussen A, Shepherd M, Nehme Z, et al. CPR-induced consciousness: a cross-sectional study of healthcare practitioners' experience. *Australas Emerg Nurs J* 2016;19:186–90.
 28. Olaussen A, Nehme Z, Shepherd M, et al. Consciousness induced during cardiopulmonary resuscitation: an observational study. *Resuscitation* 2017;113:44–50.
 29. Bernier GM. Maintenance of consciousness during closed-chest massage. *JAMA* 1962;181:446–7.
 30. Miller JB, Davie RD, Douglas DM. The efficiency of cardiac massage in ventricular fibrillation. Description of an instance of recovery of consciousness without spontaneous heart beat. *Br J Anaesth* 1961;33:22–3.
 31. McDonald GP. Code blue stories: awake and aware in the emergency department. *Hosp Physician* 2005;41:12.
 32. Yu HY, Yeh HL, Wang SS, et al. Ultra long cardiopulmonary resuscitation with intact cerebral performance for an asystolic patient with acute myocarditis. *Resuscitation* 2007;73:307–8.
 33. Bihari S, Rajajee V. Prolonged retention of awareness during cardiopulmonary resuscitation for asystolic cardiac arrest. *Neurocrit Care* 2008;9:382–6.
 34. Tobin JM, Mihm FG. A hemodynamic profile for conscious-

- ness during cardiopulmonary resuscitation. *Anesth Analg* 2009;109:1598–9.
35. Ulrichs CJ, Bottiger BW, Padosch SA. Total recall: is it ethical not to sedate people during successful resuscitation? *Resuscitation* 2014;85:e49.
 36. Fauber J. New CPR devices save lives, Medical College study finds [Internet]. *Milwaukee Journal Sentinel*; 2011 [cited 2023 Jul 30]. Available from: <https://archive.jsonline.com/news/health/114171424.html/>
 37. Greb C, Heightman AJ. Mechanical CPR helps save the day—and the patient [Internet]. *JEMS*; 2014 [cited 2023 Jul 30]. Available from: <https://www.jems.com/patient-care/mechanical-cpr-helps-save-day-and-patient/>
 38. Gwinnutt C. Awareness during resuscitation. *Resuscitation* 2015;97:e17.
 39. Oksar M, Turhanoglu S. Is it possible to maintain consciousness and spontaneous ventilation with chest compression in the early phase of cardiac arrest? *Case Rep Anesthesiol* 2016; 2016:3158015.
 40. Pound J, Verbeek PR, Cheskes S. CPR induced consciousness during out-of-hospital cardiac arrest: a case report on an emerging phenomenon. *Prehosp Emerg Care* 2017;21:252–6.
 41. Rice DT, Nudell NG, Habrat DA, Smith JE, Ernest EV. CPR induced consciousness: it's time for sedation protocols for this growing population. *Resuscitation* 2016;103:e15–6.
 42. Grandi T, De Carlo S, Carosi V, et al. Six cases of CPR-induced consciousness in witnessed cardiac arrest. *Italian J Emerg Med* 2017;12:1–4.
 43. Gray R. Consciousness with cardiopulmonary resuscitation. *Can Fam Physician* 2018;64:514–7.
 44. Wacht O, Huri R, Strugo R. Case study: combative cardiac patient: what do you do when a patient regains consciousness during mechanical CPR? *EMS World* 2015;44:29–33.
 45. Pinto J, Almeida P, Ribeiro F, Simoes R. Cardiopulmonary resuscitation induced consciousness a case report in an elderly patient. *Eur J Case Rep Intern Med* 2020;7:001409.
 46. Sukumar V. Having a conscious patient during cardiopulmonary resuscitation: is it not time to consider sedation protocol?: a case report. *A A Pract* 2019;13:250–2.
 47. Asghar A, Salim B, Tahir S, Islam F, Khan MF. Awareness during cardiopulmonary resuscitation. *Indian J Crit Care Med* 2020;24:136–7.
 48. Chin KC, Yang SC, Chiang WC. Video of cardiopulmonary resuscitation induced consciousness during ventricular fibrillation. *Resuscitation* 2020;155:22–3.
 49. Lapostolle F, Petrovic T, Alheritiere A, Agostinucci JM, Adnet F. Life signs in "dead" patients. *Resuscitation* 2012;83:e164.
 50. Al Harbi MK, Alattas KA, Alnajar M, Albuthi MF. Prone cardiopulmonary resuscitation in elderly undergoing posterior spinal fusion with laminectomy. *Saudi J Anaesth* 2020;14: 123–6.
 51. Brown J, Rogers J, Soar J. Cardiac arrest during surgery and ventilation in the prone position: a case report and systematic review. *Resuscitation* 2001;50:233–8.
 52. Bustillo MA, Lien CA, Mack PF, et al. Optimizing patient access during an emergency while using intraoperative computed tomography. *World Neurosurg* 2019;121:274–8.
 53. Dequin PF, Hazouard E, Legras A, Lanotte R, Perrotin D. Cardiopulmonary resuscitation in the prone position: Kouwenhoven revisited. *Intensive Care Med* 1996;22:1272.
 54. Dooney N. Prone CPR for transient asystole during lumbosacral spinal surgery. *Anaesth Intensive Care* 2010;38:212–3.
 55. de Souza Gomes D, Darcy Alves Bersot C. Cardiopulmonary resuscitation in the prone position. *Open J Anesthesiol* 2012; 2:199–201.
 56. Haffner E, Sostarich AM, Fasel T. Successful cardiopulmonary resuscitation in prone position. *Anaesthesist* 2010;59:1099–101.
 57. Loewenthal A, De Albuquerque AM, Lehmann-Meurice C, Otteni JC. Efficacy of external cardiac massage in a patient in the prone position. *Ann Fr Anesth Reanim* 1993;12:587–9.
 58. Mishra N, Singh S, Elayat A, Kaushal A. Cardiac arrest in the prone position caused by central venous cannulation-induced cardiac tamponade. *Korean J Anesthesiol* 2019;72:394–5.
 59. Miranda CC, Newton MC. Successful defibrillation in the prone position. *Br J Anaesth* 2001;87:937–8.
 60. Sun WZ, Huang FY, Kung KL, Fan SZ, Chen TL. Successful cardiopulmonary resuscitation of two patients in the prone position using reversed precordial compression. *Anesthesiology* 1992;77:202–4.
 61. Taylor JC, Buchanan CC, Rumball MJ. Cardiac arrest during craniotomy in prone position. *Trends Anaesth Crit Care* 2013; 3:224–6.
 62. Albin MS, Ritter RR, Pruett CE, Kalff K. Venous air embolism during lumbar laminectomy in the prone position: report of three cases. *Anesth Analg* 1991;73:346–9.
 63. Chen HL, Wong CS, Ho ST, Chang FL, Hsu CH, Wu CT. A lethal pulmonary embolism during percutaneous vertebroplasty. *Anesth Analg* 2002;95:1060–2.
 64. Dumont TM, Stockwell DW, Horgan MA. Venous air embolism: an unusual complication of atlantoaxial arthrodesis: case report. *Spine (Phila Pa 1976)* 2010;35:E1238–40.
 65. Ewah B, Calder I. Intraoperative death during lumbar discectomy. *Br J Anaesth* 1991;66:721–3.

66. Miyakoshi N, Hongo M, Kasukawa Y, Ishikawa Y, Kudo D, Shimada Y. Intraoperative visible air bubbling recorded as a sign of massive venous air embolism during prone position surgery for extensive ossification of spinal ligaments: a case report with a video clip. *World Neurosurg* 2019;131:38–42.
67. Pan Y, Qiu B, Yu F, Hu B. Fatal air embolism during endoscopic retrograde cholangio-pancreatography (ERCP): a case report. *J Med Coll PLA* 2012;27:239–43.
68. Pinheiro LC, Carmona BM, de Nazareth Chaves Fascio M, de Souza IS, de Azevedo RA, Barbosa FT. Cardiac arrest after epidural anesthesia for a esthetic plastic surgery: a case report. *Rev Bras Anesthesiol* 2017;67:544–7.
69. Mazer SP, Weisfeldt M, Bai D, et al. Reverse CPR: a pilot study of CPR in the prone position. *Resuscitation* 2003;57:279–85.
70. Wei J, Tung D, Sue SH, Wu SV, Chuang YC, Chang CY. Cardiopulmonary resuscitation in prone position: a simplified method for outpatients. *J Chin Med Assoc* 2006;69:202–6.
71. Burki AM, Mahboob S, Fatima T. CPR in prone position during neurosurgery. *Anaesth Pain Intensive Care* 2017;21:275–8.
72. Gueugniaud PY, Muchada R, Bertin-Maghit M, Griffith N, Petit P. Non-invasive continuous haemodynamic and PET-CO₂ monitoring during peroperative cardiac arrest. *Can J Anaesth* 1995;42:910–3.
73. Tofil NM, Dollar J, Zinkan L, et al. Performance of anesthesia residents during a simulated prone ventricular fibrillation arrest in an anesthetized pediatric patient. *Paediatr Anaesth* 2014;24:940–4.
74. Parshuram CS, Dryden-Palmer K, Farrell C, et al. Effect of a Pediatric Early Warning System on all-cause mortality in hospitalized pediatric patients: the EPOCH Randomized Clinical Trial. *JAMA* 2018;319:1002–12.
75. Agulnik A, Mora Robles LN, Forbes PW, et al. Improved outcomes after successful implementation of a pediatric early warning system (PEWS) in a resource-limited pediatric oncology hospital. *Cancer* 2017;123:2965–74.
76. Bonafide CP, Localio AR, Roberts KE, Nadkarni VM, Weirich CM, Keren R. Impact of rapid response system implementation on critical deterioration events in children. *JAMA Pediatr* 2014;168:25–33.
77. Brill R, Gibson R, Luria JW, et al. Implementation of a medical emergency team in a large pediatric teaching hospital prevents respiratory and cardiopulmonary arrests outside the intensive care unit. *Pediatr Crit Care Med* 2007;8:236–46.
78. Hanson CC, Randolph GD, Erickson JA, et al. A reduction in cardiac arrests and duration of clinical instability after implementation of a paediatric rapid response system. *Postgrad Med J* 2010;86:314–8.
79. Hunt EA, Zimmer KP, Rinke ML, et al. Transition from a traditional code team to a medical emergency team and categorization of cardiopulmonary arrests in a children's center. *Arch Pediatr Adolesc Med* 2008;162:117–22.
80. Kotsakis A, Lobos AT, Parshuram C, et al. Implementation of a multicenter rapid response system in pediatric academic hospitals is effective. *Pediatrics* 2011;128:72–8.
81. McKay H, Mitchell IA, Sinn K, et al. Effect of a multifaceted intervention on documentation of vital signs and staff communication regarding deteriorating paediatric patients. *J Paediatr Child Health* 2013;49:48–56.
82. Parshuram CS, Bayliss A, Reimer J, Middaugh K, Blanchard N. Implementing the Bedside Paediatric Early Warning System in a community hospital: a prospective observational study. *Paediatr Child Health* 2011;16:e18–22.
83. Sefton G, McGrath C, Tume L, Lane S, Lisboa PJ, Carrol ED. What impact did a Paediatric Early Warning System have on emergency admissions to the paediatric intensive care unit?: an observational cohort study. *Intensive Crit Care Nurs* 2015;31:91–9.
84. Sharek PJ, Parast LM, Leong K, et al. Effect of a rapid response team on hospital-wide mortality and code rates outside the ICU in a Children's Hospital. *JAMA* 2007;298:2267–74.
85. Tibballs J, Kinney S. Reduction of hospital mortality and of preventable cardiac arrest and death on introduction of a pediatric medical emergency team. *Pediatr Crit Care Med* 2009;10:306–12.
86. Seidler AL, Gyte GM, Rabe H, et al. Umbilical cord management for newborns < 34 weeks' gestation: a meta-analysis. *Pediatrics* 2021;147:e20200576.
87. Bowker L, Stewart K. Predicting unsuccessful cardiopulmonary resuscitation (CPR): a comparison of three morbidity scores. *Resuscitation* 1999;40:89–95.
88. Cho YJ, Kim YJ, Kim MY, et al. Validation of the Good Outcome Following Attempted Resuscitation (GO-FAR) score in an East Asian population. *Resuscitation* 2020;150:36–40.
89. George AL Jr, Folk BP 3rd, Crecelius PL, Campbell WB. Pre-arrest morbidity and other correlates of survival after in-hospital cardiopulmonary arrest. *Am J Med* 1989;87:28–34.
90. Guilbault RW, Ohlsson MA, Afonso AM, Ebell MH. External validation of two classification and regression tree models to predict the outcome of inpatient cardiopulmonary resuscitation. *J Intensive Care Med* 2017;32:333–8.
91. Hong SI, Kim YJ, Cho YJ, Huh JW, Hong SB, Kim WY. Predictive value of pre-arrest albumin level with GO-FAR score in pa-

- tients with in-hospital cardiac arrest. *Sci Rep* 2021;11:10631.
92. Ohlsson MA, Kennedy LM, Juhlin T, Melander O. Evaluation of pre-arrest morbidity score and prognosis after resuscitation score and other clinical variables associated with in-hospital cardiac arrest in southern Sweden. *Resuscitation* 2014;85:1370–4.
 93. Rubins JB, Kinzie SD, Rubins DM. Predicting outcomes of in-hospital cardiac arrest: retrospective US validation of the Good Outcome Following Attempted Resuscitation Score. *J Gen Intern Med* 2019;34:2530–5.
 94. Cartledge S, Bray JE, Leary M, Stub D, Finn J. A systematic review of basic life support training targeted to family members of high-risk cardiac patients. *Resuscitation* 2016;105:70–8.
 95. Cartledge S, Feldman S, Bray JE, Stub D, Finn J. Understanding patients and spouses experiences of patient education following a cardiac event and eliciting attitudes and preferences towards incorporating cardiopulmonary resuscitation training: a qualitative study. *J Adv Nurs* 2018;74:1157–69.
 96. Blewer AL, Putt ME, McGovern SK, et al. A pragmatic randomized trial of cardiopulmonary resuscitation training for families of cardiac patients before hospital discharge using a mobile application. *Resuscitation* 2020;152:28–35.
 97. Gonzalez-Salvado V, Abelairas-Gomez C, Gude F, et al. Targeting relatives: impact of a cardiac rehabilitation programme including basic life support training on their skills and attitudes. *Eur J Prev Cardiol* 2019;26:795–805.
 98. Han KS, Lee JS, Kim SJ, Lee SW. Targeted cardiopulmonary resuscitation training focused on the family members of high-risk patients at a regional medical center: a comparison between family members of high-risk and no-risk patients. *Ulus Travma Acil Cerrahi Derg* 2018;24:224–33.
 99. Kim HS, Kim HJ, Suh EE. The effect of patient-centered CPR education for family caregivers of patients with cardiovascular diseases. *J Korean Acad Nurs* 2016;46:463–74.
 100. Gordon M, Patricio M, Horne L, et al. Developments in medical education in response to the COVID-19 pandemic: a rapid BEME systematic review: BEME guide no. 63. *Med Teach* 2020;42:1202–15.
 101. Kent F, George J, Lindley J, Brock T. Virtual workshops to preserve interprofessional collaboration when physical distancing. *Med Educ* 2020;54:661–2.
 102. Tsang AC, Lee PP, Chen JY, Leung GK. From bedside to website: a neurological clinical teaching experience. *Med Educ* 2020;54:660.
 103. Feltes M, Becker J, McCall N, Mbanjumucyo G, Sivasankar S, Wang NE. Teaching how to teach in a train-the-trainer program. *J Grad Med Educ* 2019;11(4 Suppl):202–4.
 104. Kim EJ, Roh YS. Competence-based training needs assessment for basic life support instructors. *Nurs Health Sci* 2019;21:198–205.