

상하순에 동시에 발생한 선천성 이중구순 1예

아주대학교 의과대학 이비인후과학교실

정 연 훈 · 조 민 정

A Case of Bilabial Congenital Double Lips

Yun-Hoon Choung, DDS, MD and Min Jung Cho, MD

Department of Otolaryngology, Ajou University School of Medicine, Suwon, Korea

-ABSTRACT-

Congenital double lip deformity is an infrequent developmental abnormality affecting the lips, more commonly the upper lip. It resulted from an accessory fold of redundant mucous membrane inside the vermilion border of the lips. Especially, there is no previous report on the bilabial congenital double lips. We present a very rare case of double lips occurred simultaneously in both upper and lower lips. (J Clinical Otolaryngol 2004;15:282-285)

KEY WORDS : Double lip.

서 론

(congenital double lip)

(vermilion border)

1)

2)

1)2)

가

3)4)

2)

1

증 례

15

vermilion border

가

(labial frenulum)가

(Fig. 1).

: 2004 9 11

: 2004 11 23

: , 442 - 791

5

: (031) 219 - 5263 ·

: (031) 219 - 5264

E - mail : yhc@ajou.ac.kr

(upper labial frenulum)
 (tran-
 sverse elliptical)
 (Fig. 2).



Fig. 1. Clinical photograph demonstrating bilabial double lips. Bilateral redundant folds of soft tissue were noted on the mucosal surface of the upper and lower lips.



Fig. 2. Location of surgical excision boundary. Transverse elliptical excision was made over each side of redundant tissue.




Fig. 3. Pathologic findings (H & E stain, x 40). Hyperplastic mucosal glands were noted.




Fig. 4. The clinical photograph of 4 months after surgical excision shows normal appearance.

- rection of congenital double lip. J Kyung Hee Univ Med Center 1993;9:107-10.*
- 6) Kenny KF, Hreha JP, Dent CD. *Bilateral redundant mucosal tissue of the upper lip. J Am Dent Assoc 1990;120:193-4.*
 - 7) Lamster IB. *Mucosal reduction for correction of a maxillary double lip. Report of a case. Oral Surg Oral Med Oral Pathol 1983;55:457-8.*
 - 8) Beinhoff U, Piza-Katzer H. *Double lip in a patient with Ascher's syndrome. Eur J Plast Surg 1998;21:370-3.*
 - 9) Oldfield MC. *Prolapse of the lip mucosa. Br J Surg 1959;47:58-60.*
 - 10) Guerrero-Santos J, Altamirano JT. *The use of W-plasty for the correction of double-lip deformity. Plast Reconstr Surg 1967;39:478.*
 - 11) Reddy KA, Koteswara Rao. *Congenital double lip: A review of seven cases. Plast Reconstr Surg 1989;84(3):420-3.*
 - 12) Wood NK, Goaz PW. *Peripheral oral exophytic lesions. In: Wood NK, Goaz PW editors. Differential diagnosis of oral lesion. 3rd ed. St. Louis: The C.V. Mosby Company;1985. p.162-9.*
 - 13) Gupta IJ, Bhatnagar SK. *Congenital double lip. Ind J Plast Surg 1979;27:12.*
 - 14) Alkan A, Metin M. *Maxillary double lip: Report of two cases. J Oral Science 2001;43:69-72.*